arXiv:1303.6706v1 [math.NT] 26 Mar 2013

Formal Groups and Combinatorial Objects

Mohammad Sadek

Abstract

Let E be an elliptic curve defined over $\mathbb{Q}$ by one of the following Weierstrass equations

$$\begin{aligned} y^2+a_1xy+a_3y &= x^3+a_2x^2+a_4x, \\ y^2+a_3y &= x^3+a_6, \end{aligned}$$

where $a_i\in\mathbb{Z}$. If $z=-x/y$, we find the z-power series expansion of the invariant differential ω_E of E. As an application, we use Atkin-Swinnerton Dyer congruences to relate the coefficients of the expansion of ω_E to the coefficients of the modular form associated to E.

1 Introduction

Let E be an elliptic curve defined over $\mathbb{Q}$ described by the Weierstrass equation

$$y^2+a_1xy+a_3y=x^3+a_2x^2+a_4x+a_6, a_i\in\mathbb{Z}.$$

Choosing a local parameter $z=-x/y$ for E at its origin O_E, one can associate to E a power series $w(z)=\sum_{n=0}^{\infty}s_iz^i\in\mathbb{Z}[a_1,a_2,a_3,a_4,a_6][\![z]\!]$. Consequently, there are Laurent series expansions for the coordinates $x(z)$ and $y(z)$. In fact, since the pair $(x(z),y(z))$ gives a formal solution for the Weierstrass equation describing E, rational points on E can be produced by choosing $z\in\mathbb{Q}$.

Furthermore, the invariant differential ω_E of E can be expressed as a power series. The formal logarithm of E is the formal integral $\log_E(z)=\int\omega_E(z)$. The formal group associated to E over $\mathbb{Z}$ is $F(X,Y)=\log_E^{-1}(\log_E(X)+\log_E(Y))$.

Honda in [5] found an interesting link between the L-series $L(s)=\sum_{n=1}^{\infty}c_nn^{-s}$ of an elliptic curve E and the formal group associated to E. If one sets $g(x)=\sum_{n=1}^{\infty}n^{-1}c_nn^{-s}$, then $G(X,Y) = g^{-1}(g(X)+g(Y))$ is a formal group over $\mathbb{Z}$, moreover, $G(X,Y)$ is isomorphic to the formal group law $F(X,Y)$ associated to E over $\mathbb{Z}$. The isomorphism between these formal groups is made explicit to produce Atkin-Swinnerton Dyer congruence relations. These congruence relations connect the coefficients of the L-series and the coefficients of the power series expansion of the invariant differential ω_E. The modularity of elliptic curves E defined over $\mathbb{Q}$ implies the existence of a set of congruence relations between the coefficients of the modular form attached to E and the coefficients of the power series expansion of ω_E.

The above discussion indicates that explicit formulas for the coefficients of the power series of ω_E will provide us with information about formal groups, L-series and modular forms associated to elliptic curves. The only explicit description for the power series of ω_E can be found in [2] when E has a rational 2-torsion point, i.e., when E is described by a Weierstrass equation of the form $y^2 = x^3 + a_2x^2 + a_4x$. It is shown that $\omega_E = \sum_{n=0}^{\infty} P_n\left(a_2/\sqrt{\Delta}\right)\left(\sqrt{\Delta}\right)^n z^{2n}\,dz$, where P_n is the n-th Legendre polynomial and $\Delta = a_2^2 - a_4$. This enables the authors to find explicit congruence relations satisfied by Legendre polynomials using Atkin-Swinnerton Dyer congruences and sharper congruences using other techniques when E has complex multiplication.

More combinatorial quantities appear as coefficients of invariant differentials. In [9], the integers $(-1)^m \sum_k \binom{m}{k}^3$ appear as the coefficients of the holomorphic differential form of a model of a $K3$-surface. In [10], Apéry Numbers, $\sum_k \binom{n}{k}^2\binom{n+k}{k}$, which appear in Apéry's proof of the irrationality of $\zeta(2)$ and $\zeta(3)$ turn out to be the coefficients of the holomorphic differential form of a model of a $K3$-surface.

In this note, we investigate a larger family of elliptic curves. We write explicit formulas for the z-power series expansion of the invariant differentials of elliptic curves described by $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x$ and $y^2 + a_3y = x^3 + a_6$. One finds a power series solution to the functional equation satisfied by $w(z)$ and use this to find a z-power series expansion for ω_E. Several combinatorial numbers appear as coefficients of these power series. These numbers include

$$\sum_{m=\lfloor n/2\rfloor}^{n} \sum_{k=0}^{\lfloor m/2\rfloor} \sum_{r=0}^{n-m-k} \binom{m}{k}\binom{m-k}{k}\binom{m-2k}{r}\binom{k}{n-m-k-r}, \text{ and}$$

$$\sum_{k=\lfloor (n-1)/2\rfloor}^{n-1} \binom{n+k-1}{k}\binom{k}{n-k-1},$$

and hence we have families of congruence relations satisfied by the coefficients of the modular forms of these elliptic curves, and an explicit description of the formal logarithms of the formal groups associated to them. Indeed using these congruence relations together with Hasse's bound for modular coefficients, one can find the modular coefficients explicitly. For instance some vanishing results of modular coefficients are illustrated throughout the note.

It is worth mentioning that although one can find for any Weierstrass equation the power series $w(z)$, it is not clear for the author how to use it to obtain simple formulas for the invariant differential when $a_i \neq 0$ for every i. However, the two families of Weierstrass equations that we treat are broad enough to include elliptic curves with rational torsion points for example.

2 Formal groups of elliptic curves

Let E be an elliptic curve defined over the rational field $\mathbb{Q}$. Assume that E is described by the following Weierstrass equation

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6,\ a_i \in \mathbb{Z}.$$

We furthermore assume that the Weierstrass equation is globally minimal. Let A be the local ring of functions defined at the origin O_E, and $\widehat{A}$ the completion of A at its maximal ideal. Then $\widehat{A}$ is isomorphic to the power series ring $\mathbb{Q}[\![z]\!]$ where z is a parameter at the origin. This is used to express the Weierstrass coordinates as formal power series in z. More explicitly, we set

$$z = -\frac{x}{y},\ \text{and}\ w = -\frac{1}{y}$$

Now z is a parameter at the origin $(z, w) = (0, 0)$. The Weierstrass equation for E becomes

$$w = z^3 + a_1zw + a_2z^2w + a_3w^2 + a_4zw^2 + a_6w^3$$

substituting the equation into itself recursively we write w as a power series in z

$$w(z) = \sum_{i\geq 0} s_i z^i \in \mathbb{Z}[a_1, \ldots, a_6][\![z]\!]$$

where

$$\begin{aligned} s_0 &= s_1 = s_2 = 0,\ s_3 = 1,\ \text{ and,} \\ s_n &= a_1s_{n-1} + a_2s_{n-2} + a_3\sum_{k+l=n} s_ks_l + a_4\sum_{k+l=n-1} s_ks_l + a_6\sum_{k+l+m=n} s_ks_ls_m,\ n \geq 4. \end{aligned} \tag{1}$$

The reader can consult [1] and Chapter IV in [8]. Formal series expressions for x and y can be deduced as $x(z) = \dfrac{z}{w(z)}$ and $y(z) = \dfrac{-1}{w(z)}$. Therefore, the pair $(x(z), y(z))$ is a formal solution to the Weierstrass equation of E.

The invariant differential ω_E of E can be expressed as a formal power series in z

$$\omega_E = \frac{dy}{3x^2 + 2a_2x + a_4 - a_1y} = \sum_{n=1}^{\infty} b(n)z^{n-1}\,dz,\ b(n) \in \mathbb{Z},\ b(1) = 1.$$

The formal logarithm associated to E is $f(T) = \displaystyle\int_0^T \omega_E(T)$, hence the formal group associated to E over $\mathbb{Z}$ is given by

$$F(X, Y) = f^{-1}\left(f(X) + f(Y)\right).$$

In [5], Honda introduced an intriguing relation between the L-series of an elliptic curve and the formal group corresponding to this elliptic curve. Briefly, the local L-series L_p attached to E at some prime p is defined as follows

$$L_p(s) = (1 - t_p\,p^{-s} + u_p\,p^{1-2s})^{-1}$$

where $(t_p, u_p) = (1 + p - A_p, 1)$ if E has good reduction modulo the prime p and A_p is the number of points on the reduced curve E mod p, $(t_p, u_p) = (0, 0)$ if E has additive reduction, and $(t_p, u_p) = (1, 0)$ or $(-1, 0)$ if E has multiplicative reduction and the tangents at the double point have rational slopes over $\mathbb{F}_p$ or not respectively. The L-series associated to E is defined as

$$L(s) = \prod_p L_p(s) = \sum_{n=1}^{\infty} c_n n^{-s}$$

where $c_p = t_p$ for p prime. Let $g(x) = \displaystyle\sum_{n=1}^{\infty} n^{-1}c_nx^n$. The formal group associated to the L-series is defined as $G(x, y) = g^{-1}(g(x) + g(y))$. The following theorem was proved by Honda in [5] for elliptic curves with no bad primes at 2 and 3, and was extended in [4] to include any elliptic curve over $\mathbb{Q}$.

Theorem 2.1. *Let E be an elliptic curve over $\mathbb{Q}$ and let $F(x, y)$ be the formal group associated to E over $\mathbb{Z}$. Let $G(x, y)$ be the formal group associated to the L-series of E. Then $F(x, y)$ is defined over $\mathbb{Z}$ and is isomorphic over $\mathbb{Z}$ to $G(x, y)$, i.e., there exists a power series $\phi(X) \in \mathbb{Z}[\![X]\!]$ such that $\phi(F(X, Y)) = G(\phi(X), \phi(Y))$.*

The above theorem implies that $g(x) = f(\phi^{-1}(X))$. Expanding this out in terms of coefficients yields the Atkin-Swinnerton Dyer (Honda-Cartier) congruences, see [3].

Theorem 2.2. *Let* $E, b(n), t_p, u_p$ *be defined as above. One has*

$$b(np) - t_p\, b(n) + p\, u_p\, b(n||p) \equiv 0 \ mod\ p^s \ \ if\ n \equiv 0 \ mod\ p^{s-1}$$

where $b(n||p) = 0$ *if* $p \nmid n$ *and* $b(n||p) = b(n/p)$ *if* $p \mid n$.

Corollary 2.3. *If* E *has good reduction mod* p*, then*

a) $b(p) \equiv t_p \ mod\ p$;

b) $b(np) \equiv b(n)\, b(p) \ mod\ p \ if\ p \nmid n$;

c) $b(np) - t_p\, b(n) + p\, b(n/p) \equiv 0 \ mod\ p^s \ if\ n \equiv 0 \ mod\ p^{s-1}, s \geq 1$.

If E *has multiplicative reduction, then*

a) $b(p) \equiv t_p \ mod\ p$;

b) $b(np) \equiv t_p\, b(n) \ mod\ p^s \ for\ n \equiv 0 \ mod\ p^{s-1}, s \geq 1$.

where $t_p = 1$ *or* -1 *according to the tangents at the double point having rational slopes over* $\mathbb{F}_p$ *or not respectively.*
If E *has additive reduction mod* p*, then* $b(np) \equiv 0 \ mod\ p^s \ if\ n \equiv 0 \ mod\ p^{s-1}$.

The modulartiy of elliptic curves over $\mathbb{Q}$ implies that if the L-series of E is $L_E(s) = \sum_{n=1}^{\infty} c_n n^{-s}$, then the generating function $f(q) = \sum_{n=1}^{\infty} c_n q^n$ of the coefficients c_n is a modular form where $q = e^{2\pi i \tau}$.

As we will compute different power series expansions, we will need Lagrange Inversion Theorem. If $\phi(x)$ is a power series, then $[x^n]\phi(x)$ denotes the coefficient of x^n in $\phi(x)$.

Lemma 2.4. *Suppose* $u = u(x)$ *is a power series in* x *satisfying* $x = u/\phi(u)$ *where* $\phi(u)$ *is a power series in* u *with a nonzero constant term. Then we have*

$$[x^n]u(x) = \frac{1}{n}[u^{n-1}]\phi^n(u).$$

3 The elliptic curve $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x$

In this section we consider elliptic curves over $\mathbb{Q}$ given by the Weierstrass equation $y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x$ where $a_i \in \mathbb{Z}$. Set $z = -x/y$ to be a parameter at the origin, and $w=-1/y$. According to eq (1), $w(z)=\sum_{n=0}^{\infty} s_n z^n$, where $s_0=s_1=s_2=0, s_3=1$ and

$$s_n = a_1 s_{n-1} + a_2 s_{n-2} + a_3 \sum_{k+l=n} s_k s_l + a_4 \sum_{k+l=n-1} s_k s_l.$$

The generating function $w(z)$ of the sequence $(s_n)_{n=0}^{\infty}$ satisfies the following functional equation:

$$w(z) = z^3 + a_1 z w(z) + a_2 z^2 w(z) + a_3 w(z)^2 + a_4 z w(z)^2.$$

The above equation is quadratic in $\omega(z)$. As a consequence, one has

$$w(z) = \frac{1 - a_1 z - a_2 z^2 - \sqrt{(1 - a_1 z - a_2 z^2)^2 - 4z^3(a_3 + a_4 z)}}{2(a_3 + a_4 z)}. \tag{2}$$

We chose the negative sign because the positive sign would force $w(z)$ to have a pole at $z=0$. Recall that $x(z)=z/w(z)$ and $y(z)=-1/w(z)$. The invariant differential ω_E of E is given by

$$\begin{aligned} \omega_E &= \frac{dy}{3x^2 + 2a_2 x + a_4 - a_1 y} = \frac{\frac{d}{dz}(-1/w)}{3(z/w)^2 + 2a_2(z/w) + a_4 - a_1(-1/w)}\, dz \\ &= \frac{\frac{d}{dz}\,\omega}{3z^2 + 2a_2 z w + a_4 w^2 + a_1 w}\, dz. \end{aligned} \tag{3}$$

We use Mathematica, [7], to substitute (2) in the formula for ω_E to obtain

$$\begin{aligned} \omega_E &= \frac{1}{\sqrt{(1 - a_1 z - a_2 z^2)^2 - 4z^3(a_3 + a_4 z)}}\, dz \\ &= \frac{1}{\sqrt{1 - 2(a_1 + a_2 z)z + \left[(a_1 + a_2 z)^2 - 4(a_3 + a_4 z)z\right] z^2}}\, dz. \end{aligned}$$

Before we proceed to the main result of this section we need to recall the definitions of Legendre polynomials $P_n(x)$ and generalized central trinomial polynomials $T_n(x,y)$, see [11] for example.

A Legendre polynomial $P_n(x)$, $n \ge 0$, is defined as follows

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n}(x^2-1)^n = \frac{1}{2^n} \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n}{k} \binom{2n-2k}{n} x^{n-2k},$$

and the generating function of Legendre polynomials is given by

$$\frac{1}{\sqrt{1-2xt+t^2}} = \sum_{n=0}^{\infty} P_n(x) t^n.$$

The generalized central trinomial polynomial $T_n(x,y)$ is given as follows

$$T_n(x,y) = [t^n](t^2+xt+y)^n = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n}{k}\binom{n-k}{k} x^{n-2k}y^k,$$

and the generating function is given by

$$\frac{1}{\sqrt{1-2xt+(x^2-4y)t^2}} = \sum_{n=0}^{\infty} T_n(x,y)t^n.$$

One can deduce an immediate relation between the polynomials T_n and P_n, namely

$$T_n(x,y) = (\sqrt{x^2-4y})^n P_n\left(\frac{x}{\sqrt{x^2-4y}}\right).$$

Theorem 3.1. *Let $E: y^2+a_1xy+a_3y = x^3+a_2x^2+a_4x,\ a_i \in \mathbb{Z}$, be an elliptic curve over $\mathbb{Q}$. Let $z = -x/y$ be a local parameter at O_E. The z-power series expansion of ω_E is given by $\sum_{n=0}^{\infty} b(n+1)\, z^n\, dz$, where $b(n+1)$ is*

$$\sum_{m=\lfloor n/2 \rfloor}^{n} \sum_{k=0}^{\lfloor m/2 \rfloor} \sum_{r=0}^{n-m-k} \binom{m}{k}\binom{m-k}{k}\binom{m-2k}{r}\binom{k}{n-m-k-r} a_1^{m-2k-r} a_2^r a_3^{2k-n+m+r} a_4^{n-m-k-r}$$

Proof: We have already shown that $\omega_E = \dfrac{1}{\sqrt{1-2(a_1+a_2z)z+[(a_1+a_2z)^2-4(a_3+a_4z)z]\,z^2}}\,dz.$ We compare the formula for ω_E with the generating function of the generalized central trinomial polynomials. One concludes that

$$\omega_E \;=\; \sum_{m=0}^{\infty} T_m\left(a_1+a_2z,(a_3+a_4z)z\right)\, z^m\, dz.$$

Now one has

$$\begin{aligned}
T_m(a_1+a_2z,(a_3+a_4z)z) &= \sum_{k=0}^{\lfloor m/2 \rfloor} \binom{m}{k}\binom{m-k}{k}(a_1+a_2z)^{m-2k}(a_3+a_4z)^k z^k \\
&= \sum_{k=0}^{\lfloor m/2 \rfloor} \binom{m}{k}\binom{m-k}{k}\left(\sum_{i=0}^{m-2k}\binom{m-2k}{i} a_1^{m-2k-i}a_2^i\, z^i\right)\left(\sum_{j=0}^{k}\binom{k}{j} a_3^{k-j}a_4^j z^j\right) z^k \\
&= \sum_{k=0}^{\lfloor m/2 \rfloor} \binom{m}{k}\binom{m-k}{k}\sum_{s=0}^{m-k}\left(\sum_{r=0}^{s}\binom{m-2k}{r}\binom{k}{s-r} a_1^{m-2k-r}a_2^r a_3^{k-s+r}a_4^{s-r}\right) z^{s+k}
\end{aligned}$$

For the third equality we applied the formula for multiplying polynomials. Now,

$$\begin{aligned} b(n+1) &= [z^n]\,\omega_E = [z^n]\sum_{m=0}^{\infty} T_m\left(a_1+a_2z,(a_3+a_4z)z\right)\, z^m \\ &= \quad [z^n]\sum_{m=0}^{\infty}\sum_{k=0}^{\lfloor m/2\rfloor}\sum_{s=0}^{m-k}\sum_{r=0}^{s}\binom{m}{k}\binom{m-k}{k}\binom{m-2k}{r}\binom{k}{s-r}a_1^{m-2k-r}a_2^{r}a_3^{k-s+r}a_4^{s-r}z^{s+k+m} \end{aligned}$$

Therefore, $b(n+1)$ consists of the sum of the coefficients for which $s+k+m=n$. Since $0\le s\le m-k$, one has $n\le 2m$. Moreover, m cannot exceed n. It follow that $b(n+1)$ is

$$\sum_{m=\lfloor n/2\rfloor}^{n}\sum_{k=0}^{\lfloor m/2\rfloor}\sum_{r=0}^{n-m-k}\binom{m}{k}\binom{m-k}{k}\binom{m-2k}{r}\binom{k}{n-m-k-r}a_1^{m-2k-r}a_2^{r}a_3^{2k-n+m+r}a_4^{n-m-k-r}.$$

□

The following corollary presents a set of congruence relations satisfied by $b(n+1)$ modulo all but finitely many primes.

Corollary 3.2. *Let $E/\mathbb{Q}$ be an elliptic curve described by $y^2+a_1xy+a_3y=x^3+a_2x^2+a_4x$. Let Δ_E be the minimal discriminant of E. Let p be a prime satisfying $p\nmid\Delta_E$, A_p the number of rational points on E mod p, and $t_p=1+p-A_p$. Assume further that $\omega_E=\sum_{n=1}^{\infty}b(n)\,z^{n-1}\,dz$ where $z=-x/y$ and $b(n)$ is given in Theorem 3.1. Then*

- *a) $b(p)\equiv t_p$ mod p;*
- *b) $b(np)\equiv b(n)\,b(p)$ mod p if $p\nmid n$;*
- *c) $b(np)-t_p\,b(n)+p\,b(n/p)\equiv 0$ mod p^s if $n\equiv 0$ mod $p^{s-1}, s\ge 1$.*

Proof: This is corollary 2.3. □

We will give a few examples of families of elliptic curves E with interesting z-power series expansion of ω_E.

Example 3.3. Let $E/\mathbb{Q}$ be an elliptic curve described by $y^2=x^3+a_4x$. One has

$$\omega_E=\frac{1}{\sqrt{1-4a_4z^4}}\,dz=\sum_{n=0}^{\infty}\binom{2n}{n}a_4^n z^{(4n+1)-1}\,dz.$$

Similarly, the invariant differential of $E: y^2+a_3y=x^3$ is

$$\omega_E=\frac{1}{\sqrt{1-4a_3z^3}}\,dz=\sum_{n=0}^{\infty}\binom{2n}{n}a_3^n z^{(3n+1)-1}\,dz.$$

Corollary 3.4. *Let a be a nonzero integer, and p a prime such that $p \nmid 2a$. Set $t_p = 1 + p - A_p$ where A_p is the number of points on $E : y^2 = x^3 + ax$ mod p. One has*

a) $t_p = 0$ *if* $p \equiv 3$ *mod* 4;

If $p \equiv 1$ *mod* 4, *then*

b) $t_p \equiv \binom{\frac{p-1}{2}}{\frac{p-1}{4}}$ *mod* p;

c) $\binom{\frac{np-1}{2}}{\frac{np-1}{4}} \equiv \binom{\frac{n-1}{2}}{\frac{n-1}{4}}\binom{\frac{p-1}{2}}{\frac{p-1}{4}}$ *mod* p *if* $n \equiv 1$ *mod* 4.

d) $\binom{\frac{p^2-1}{2}}{\frac{p^2-1}{4}} - t_p\binom{\frac{p-1}{2}}{\frac{p-1}{4}} + p \equiv 0$ *mod* p^2.

Proof: One needs to observe that $b(n) = 0$ if $n \equiv 3$ mod 4, and that $\Delta_E = -64a^3$. For (a), a theorem of Hasse implies that $|t_p| < 2\sqrt{p}$. Since $t_p \equiv 0$ mod p if $p \equiv 3$ mod 4, it follows that $t_p = 0$. □

Corollary 3.5. *Let a be a nonzero integer, and p a prime such that $p \nmid 3a$. Set $t_p = 1 + p - A_p$ where A_p is the number of points on $E : y^2 + ay = x^3$ mod p. One has*

a) $t_p = 0$ *if* $p \equiv 2$ *mod* 3;

If $p \equiv 1$ *mod* 3, *then*

b) $t_p \equiv \binom{2\left(\frac{p-1}{3}\right)}{\frac{p-1}{3}}$ *mod* p;

c) $\binom{2\left(\frac{np-1}{3}\right)}{\frac{np-1}{3}} \equiv \binom{2\left(\frac{n-1}{3}\right)}{\frac{n-1}{3}}\binom{2\left(\frac{p-1}{3}\right)}{\frac{p-1}{3}}$ *mod* p *if* $n \equiv 1$ *mod* 3.

d) $\binom{2\left(\frac{np-1}{3}\right)}{\frac{np-1}{3}} - t_p\binom{2\left(\frac{n-1}{3}\right)}{\frac{n-1}{3}} + p\binom{2\left(\frac{n/p-1}{3}\right)}{\frac{n/p-1}{3}} \equiv 0$ *mod* p^s, *if* $n \equiv p^{s-1}$, $s \geq 1$. *In particular,* $\binom{2\left(\frac{p^2-1}{3}\right)}{\frac{p^2-1}{3}} - t_p\binom{2\left(\frac{p-1}{3}\right)}{\frac{p-1}{3}} + p \equiv 0$ *mod* p^2.

Proof: The corollary follows from $b(n) = 0$ if $n \equiv 2$ mod 3, and $\Delta_E = -27a^4$. □

The following example appears in the proof of Theorem 1 of [2].

Example 3.6. The invariant differential of $E : y^2 = x^3 + a_2x^2 + a_4x$ is given by

$$\omega_E = \frac{1}{\sqrt{1 - 2a_2z^2 + (a_2^2 - 4a_4)z^4}} = \sum_{n=0}^{\infty} T_n(a_2, a_4)z^{2n}.$$

We can now relate $T_n(a_2, a_4)$ to the coefficients of the modular form attached to E using Corollary 2.3.

4 The elliptic curve $y^2 + a_3y = x^3 + a_6$

We consider the family of elliptic curves defined by $y^2 + a_3y = x^3 + a_6$, $a_i \in \mathbb{Z}$. We set $z = -x/y$ and $w(z) = -1/y = \sum_{n=0}^{\infty} s_nz^n$, where $s_0 = s_1 = s_2 = 0, s_3 = 1$ and $s_n = a_3 \sum_{k+l=n} s_ks_l + a_6 \sum_{k+l+m=n} s_ks_ls_m$, $n \geq 4$. Now $w(z)$ satisfies the following functional equation

$$w(z) = z^3 + a_3w(z)^2 + a_6w(z)^3.$$

The latter equation can be written as

$$v = z^3 = w(z)\left(1 - a_3w(z) - a_6w(z)^2\right) = \frac{w(z)}{1/\left(1 - a_3w(z) - a_6w(z)^2\right)}.$$

We now can apply Lagrange inversion Theorem, Lemma 2.4. One obtains

$$\begin{aligned}
[v^n]w(v) &= \frac{1}{n}[w^{n-1}]\left(\frac{1}{1 - a_3w - a_6w^2}\right)^n \\
&= \frac{1}{n}[w^{n-1}]\sum_{k=0}^{\infty}(-1)^k\binom{-n}{k}(a_3 + a_6w)^kw^k \\
&= \frac{1}{n}[w^{n-1}]\sum_{k=0}^{\infty}\sum_{j=0}^{k}\binom{n+k-1}{k}\binom{k}{j}a_3^{k-j}a_6^jw^{j+k} \\
&= \frac{1}{n}\sum_{k=\lfloor(n-1)/2\rfloor}^{n-1}\binom{n+k-1}{k}\binom{k}{n-k-1}a_3^{2k-n+1}a_6^{n-1-k}.
\end{aligned}$$

Theorem 4.1. *Let E be an elliptic curve defined over $\mathbb{Q}$ by the Weierstrass equation $y^2 + a_3y = x^3 + a_6$. Let $z = -x/y$. The z-power series expansion of the invariant differential ω_E of E is given by*

$$\omega_E = \sum_{n=1}^{\infty}\left(\sum_{k=\lfloor(n-1)/2\rfloor}^{n-1}\binom{n+k-1}{k}\binom{k}{n-k-1}a_3^{2k-n+1}a_6^{n-k-1}\right)z^{(3n-2)-1}\,dz.$$

Proof: The invariant differential of E is defined by $\omega_E = \frac{dw/dz}{3z^2}\,dz$, see (3). The result now follows as we have shown that

$$w(z) = \sum_{n=1}^{\infty}\left(\frac{1}{n}\sum_{k=\lfloor (n-1)/2\rfloor}^{n-1}\binom{n+k-1}{k}\binom{k}{n-k-1}a_3^{2k-n+1}a_6^{n-k-1}\right)z^{3n}.$$

□

One can apply Atkin-Swinnerton Dyer congruences to the elliptic curve E defined over $\mathbb{Q}$ by the Weierstrass equation $y^2 + a_3 y = x^3 + a_6$. In particular, if p is a good prime of E, then using Hasse's bound, $|t_p| < 2\sqrt{p}$, one has $t_p = 0$ if $p \equiv 2 \bmod 3$, moreover, $t_p \equiv \sum_{k=\lfloor (p-1)/6\rfloor}^{(p-1)/3}\binom{\frac{p-1}{3}+k}{k}\binom{k}{\frac{p-1}{3}-k}a_3^{2k-\frac{p-1}{3}}a_6^{\frac{p-1}{3}-k} \bmod p$ if $p \equiv 1 \bmod 3$, where $t_p = 1 + p - A_p$ and A_p is the number of points on E mod p.

The following example corrects the recurrence relation in Theorem 1.2 of [6].

Example 4.2. The invariant differential of the elliptic curve $E : y^2 = x^3 + a_6$ is

$$\omega_E = \sum_{k=1}^{\infty}\binom{3k-3}{k-1}a_6^{k-1}z^{(6k-5)-1}\,dz.$$

Therefore, $t_p = 0$ if $p \not\equiv 1 \bmod 6$, and $t_p \equiv \binom{\frac{p-1}{2}}{\frac{p-1}{6}}a_6^{(p-1)/6}$ if $p \equiv 1 \bmod 6$.

Remark 4.3. Given Theorem 3.1 and Theorem 4.1, one can produce a large number of congruence relations relating combinatorial objects to coefficients of modular forms, and congruences satisfied by the combinatorial quantities themselves. All one needs is applying these theorems and Atkin-Swinnerton Dyer congruences to different families of elliptic curves described by Weierstrass equations of the form given in §3 and §4.

For example, elliptic curves with n-torsion points are parametrized by Tate's normal form. In other words, an elliptic curve $E_n/\mathbb{Q}$ with a rational n-torsion point $(0,0)$, $n \geq 4$, is described by the following Weierstrass equation

$$y^2 + (1-c)xy - by = x^3 - bx^2,\ c, b \in \mathbb{Z}.$$

According to Theorem 3.1, the invariant differential of E_n is

$$\omega_{E_n} = \frac{dz}{\sqrt{1 - 2(1-c-bz)z + [(1-c-bz)^2 + 4bz]z^2}} = \sum_{n=1}^{\infty} b(n)z^{n-1}\,dz,$$

where

$$b(n) = \sum_{m=0}^{n-1} \sum_{k=0}^{\lfloor m/2 \rfloor} \binom{m}{k} \binom{m-k}{k} \binom{m-2k}{n-m-k-1} (1-c)^{2m-n-k+1} (-b)^{n-m-1}.$$

One obtains congruence relations satisfied by $b(n)$ using Corollary 3.2. For instance, if we assumed further that $c = b = 1$, then the L-function of E is given by

$$L(s) = \sum_{n=1}^{\infty} \frac{c_n}{n^s}, \text{ where } \sum_{n=1}^{\infty} c_n q^n = q \prod_{n=1}^{\infty} (1-q^n)^2 \prod_{n=1}^{\infty} (1-q^{11n})^2,$$

see [12]. Therefore, if $p \neq 11$, then one gets

$$b(np) - c_p\, b(n) + p\, b(n/p) \equiv 0 \bmod p^s \text{ if } n \equiv 0 \bmod p^{s-1}, s \geq 1,$$

where

$$b(n) = \sum_{m=0}^{n-1} \sum_{k=0}^{\lfloor m/2 \rfloor} (-1)^{n-m-1} \binom{m}{k} \binom{m-k}{k} \binom{m-2k}{n-m-k-1}.$$

References

[1] A. Bluher. A leisurely introduction to formal groups and elliptic curves. http://www.math.uiuc.edu/Algebraic-Number-Theory/0076/.

[2] M. J. Coster and L. Van Hamme. Supercongruences of Atkin and Swinnerton-Dyer type for Legendre polynomials. *Journal of Number Theory*, 38:265–286, 1991.

[3] M. Hazewinkel. Three lectures on formal groups. *Canadian Mathematical Society Conference Proceedings*, 5:51–67, 1986.

[4] W. Hill. Formal groups and zeta-functions of elliptic curves. *Inventiones Math.*, 12:321–336, 1971.

[5] T. Honda. Formal groups and zeta-functions. *Osaka, J. Math.*, 5(2):199–213, 1968.

[6] P.-R. Loh and R. C. Rhoades. p-adic and combinatorial properties of modular form coefficients. *Int. J. Number Theory*, 2:305–328, 2006.

[7] Wolfram Research. *Mathematica Edition: Version 5.0.* Wolfram Research.

[8] J. Silverman. *The arithmetic of elliptic curves.* GTM 106. Springer-Verlag, New York, 1986.

[9] J. Stienstra. Formal groups and congruences for L-functions. *American Journal of Mathematics*, 109(6):1111–1127, 1987.

[10] J. Stienstra and F. Beukers. On the Picard-Fuchs equation and the formal Brauer group of certain elliptic $K3$-surfaces. *Math. Annalen*, 271:269–304, 1985.

[11] Z. W. Sun. Congruences involving generalized central trinomial coefficients. 2011. preprint, available at `http://arxiv.org/pdf/1008.3887v11.pdf`.

[12] J. Tate. The arithmetic of elliptic curves. *Inventiones Math.*, 23:179–206, 1974.

Department of Mathematics and Actuarial Science
American University in Cairo
mmsadek@aucegypt.edu